



**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



PCT/EP 03/06 168

10/517646

10 DEC 2004 INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 17 JUL 2003

WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.


In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.


Signed

Dated 29 April 2003



Patents Form 1/77

Patents Act 1977
(Rule 16)



**The
Patent
Office**

14JUN02 E725658-1 D00019
P01/7700 0.00-0213598.6

Request for grant of a patent

(See the notes on the back of this form. You can also
get an explanatory leaflet from the Patent Office to
help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

113 JUN 2002

1. Your reference

P030893GB

2. Patent application number

(The Patent Office will fill in this part)

0213598.6

3. Full name, address and postcode of the or of
each applicant (underline all surnames)

Bayer AG
D-51368 Leverkusen
Germany

8100414001

Patents ADP number (if you know it)

If the applicant is a corporate body, give the
country/state of its incorporation

Germany

4. Title of the invention

Carboxamides derivatives

5. Name of your agent (if you have one)

Carpmaels & Ransford

"Address for service" in the United Kingdom
to which all correspondence should be sent
(including the postcode)

43 Bloomsbury Square
London
WC1A 2RA

Patents ADP number (if you know it)

83001 ✓

6. If you are declaring priority from one or more
earlier patent applications, give the country
and the date of filing of the or of each of these
earlier applications and (if you know it) the or
each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise
derived from an earlier UK application,
give the number and the filing date of
the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right
to grant of a patent required in support of
this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an
applicant, or
- c) any named applicant is a corporate body

Yes

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	29
Claim(s)	5
Abstract	1
Drawing(s)	0

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)	1
--	---

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature Carpmaels & Ransford Date 13th June 2002

Carpmaels & Ransford

12. Name and daytime telephone number of person to contact in the United Kingdom Peter M. Johnston 020-7242 8692

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.

Carboxamides derivatives**Detailed Description of Invention**

5

Technical Field

10

The present invention relates to a carboxamides which are useful as an active ingredient of pharmaceutical preparations. The carboxamides of the present invention have PGI₂ [prostaglandin I₂, prostacyclin] antagonistic activity, and can be used for the prophylaxis and treatment of diseases associated with PGI₂ activity.

15

More specifically, the carboxamides derivatives of the present invention are useful for treatment and prophylaxis of urological diseases or disorders.

20

BACKGROUND ART

25

30

Prostaglandins (or prostanoids, PGs) are a group of bioactive lipid mediators generated from membrane phospholipids. They are formed from 20-carbon essential fatty acids containing 3, 4, or 5 double bonds, and carry a cyclopentane ring. They are divided into 6 main classes (D, E, F, G, H or I) by the cyclopentane ring structure. The main classes are further subdivided by subscripts 1, 2, or 3, reflecting their fatty acid precursors. PGI₂ is a member of prostanoids, and it has a double ring structure and is derived from arachidonic acid. The receptor for PGI₂ is a seven transmembrane G-protein coupled receptor, called IP. IP couples at least to G_s-type G-protein, and activates adenylyl cyclase and phospholipase C. The expression of IP

is demonstrated in aorta, coronary/pulmonary/cerebral arteries, platelets, lung, and dorsal root ganglions in addition to several other tissues.

5 One of the well-known actions of PGI₂ is for blood vessels to cause vasodilation and hypotension. Especially in septic shock, PGI₂ is produced and participate in the induction of systemic hypotension (G.D. Bottoms et al, Am J Vet Res 1982, 43(6), 999-1002). Therefore, IP receptor antagonists may prevent hypotension associated with septic shock.

10 Another well-known action of PGI₂ is for platelets to suppress aggregation. In the IP receptor knock out mice, FeCl₃-induced thrombosis formation was enhanced compared to that in wild type mice (T. Murata et al, Nature 1997, 388, 678- 682.), confirming the involvement of IP receptor in the platelet inhibition. Therefore, IP receptor antagonists may enhance the platelet activation and suppress excessive
15 bleeding such as, but not limited to, hemophilia and hemorrhage.

PGI₂ also participate in the inflammation. In the inflamed tissue, various inflammatory mediators, including prostaglandins, are produced. PGI₂ is also generated and induces vasodilation to increase blood flow. This enhances vascular
20 permeability, edema formation and leukocyte inflammation in the inflamed region (T. Murata et al, Nature 1997, 388, 678- 682.). Therefore, PGI₂ receptor antagonists may be efficacious for the treatment of inflammation.

25 PGI₂ may be involved in the pathogenesis of respiratory allergy or asthma. It is spontaneously generated and the major prostaglandin in human lung, and the appropriate antigen challenge increases PGI₂ production (E.S. Schulman et al, J Appl Physiol 1982, 53(3), 589-595.). Therefore, IP antagonists may have a utility for the treatment of those respiratory diseases.

30 In addition, an important role of IP receptor in the induction of hyperalgesia has been clearly shown by IP receptor knockout mice (T. Murata et al., Nature 1997, 388,


678-682.). Injection of acetic acid into the peritoneal cavity induced production of PGI₂. This PGI₂ is considered to bind to IP receptor on sensory neurons. As IP receptor couples to the activation of both adenylate cyclase and phospholipase C, cAMP-dependent protein kinase (PKA) and protein kinase C (PKC) are activated. PKA and PKC are known to modulate ion channels on sensory neurons such as VR1, P2X₃, and TTX-R. As a result, PGI₂ sensitizes sensory neurons to enhance the release of neurotransmitters. Hence, acetic acid injection induces nociceptive response (writhing) in mice. This acetic acid-induced writhing was greatly reduced in PGI₂ receptor-null mice as the same level as indomethacin-treated wild type mice. Several other in vivo hyperalgesia studies in rodents and in vitro studies further support that PGI₂ plays a major role in the induction of hyperalgesia and that PGI₂ acts as important modulator of sensory neurons (K. Bley et al, Trends in Pharmacological Sciences 1998, 19(4), 141-147.). Therefore, PGI₂ receptor antagonists may be useful for the treatment of pain.

Sensory neurons play very important roles not only in the pain sensation but also in the sensation of bladder distension. In normal subjects, A-delta sensory fibers are considered to play a major role to sense the bladder distention. However, in disease conditions of overactive bladder by, but not limited to, spinal cord injury, cystitis, Parkinson's disease, multiple sclerosis, previous cerebrovascular accident, and bladder outlet obstruction (BOO) caused by benign prostate hyperplasia (BPH), the sensitivity of C-fiber sensory neurons is upregulated and they contribute to the induction of the lower urinary tract symptoms. Treatment of overactive bladder patients with intravesical injection of capsaicin or its potent analog, resiniferatoxin, both of which desensitize VR1-positive C-fiber afferent neurons innervating the bladder, has been shown to be efficacious in several clinical trials (C. Silva et al, Eur Urol. 2000, 38(4), 444-452.). Therefore, C-fiber sensory neurons play an important role in the pathology of overactive bladder. PGI₂ is generated locally in the bladder and it is the major prostaglandin released from the human bladder. In a rabbit BOO model, a stable metabolite of PGI₂ was reported to be increased in BOO bladder (JM. Masick et al, Prostaglandins Other Lipid Mediat. 2001, 66(3), 211-219.).

Hence, PGI₂ from disease bladder sensitizes C-fiber sensory neurons, and as a result, it may induce symptoms of overactive bladder. Therefore, antagonists of PGI₂ receptor are expected to be useful in the treatment of overactive bladder and related urinary disorders.

5

EP-A-407 200 discloses antihyperlipidemics agents represented by the general formula:




10

wherein R¹' and R²' are defined in the application.

DE-A-2328391 discloses compounds that are useful for the treatment of heart diseases represented by the general formula:

15



However, none of the references and other reference discloses carboxamides derivatives having PGI₂ antagonistic activity.


20

The development of a compound which has effective PGI₂ antagonistic activity and can be used for the prophylaxis and treatment of diseases associated with PGI₂ activity has been desired.

Summary of the invention

As the result of extensive studies on chemical modification of carboxamides derivatives, the present inventors have found that the compounds of the structure related to the present invention have unexpectedly excellent PGI₂ and/or antagonistic activity. The present invention has been accomplished based on these findings.

This invention is to provide a novel carboxamide derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof:



wherein m and n independently represent an integer from 0 to 2;

-X- represents $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$, or $-\text{C}\equiv\text{C}-$;

-R¹ represents $-\text{OR}^{11}$, $-\text{SR}^{11}$, $-\text{SOR}^{11}$, $-\text{SO}_2\text{R}^{11}$, $-\text{NR}^{12}\text{R}^{13}$, or $-\text{CHR}^{14}\text{R}^{15}$,

wherein R¹¹ represents (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl,

R¹² and R¹³ independently represent hydrogen, (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl, or R¹² and R¹³ together with the nitrogen atom to which they are attached, form a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

R¹⁴ and R¹⁵ independently represent hydrogen, (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or

heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkoxy optionally substituted by aryl or heteroaryl, or

5 R¹⁴ and R¹⁵ together with the CH to which they are attached, form a (C₃₋₈)cycloalkyl optionally interrupted by NH, or O, or R¹⁴ and R¹⁵ together with the CH to which they are attached, form a phenyl optionally substituted by hydroxy, halogen or (C₁₋₆) alkyl;

10 R² represents hydrogen, cyano, (C₁₋₆) alkoxy, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, (C₃₋₇)cycloalkyl, or (C₁₋₆) alkyl optionally substituted by amino, (C₁₋₆)alkylamino, or phenyl.

15 The compounds of the present invention surprisingly show excellent PGI₂ antagonistic activity. They are, therefore, suitable for the production of medicament or medical composition, which may be useful to treat PGI₂ related diseases.


20 More specifically, since the carboxamides derivatives of the present invention antagonize PGI₂, they are useful for treatment and prophylaxis of urological diseases or disorder.

25 The compounds of the present invention are also useful for treatment of urological diseases or disorders. Such diseases or disorders include bladder outlet obstruction, overactive bladder, urinary incontinence, detrusor hyper-reflexia, detrusor instability, reduced bladder capacity, frequency of micturition, urge incontinence, stress incontinence, bladder hyperreactivity, benign prostatic hypertrophy (BPH), prostatitis, urinary frequency, nocturia, urinary urgency, pelvic hypersensitivity, urethritis, pelvic pain syndrome, prostatodynia, cystitis, or idiopathic bladder hypersensitivity.

30 The compounds of the present invention are also useful for treatment of pain including, but not limited to inflammatory pain, neuropathic pain, acute pain,

chronic pain, dental pain, premenstrual pain, visceral pain, headaches, and the like; hypotension; hemophilia and hemorrhage; inflammation; respiratory states from allergies or asthma, since the diseases also relate to PGI₂.

- 5 In another embodiment, the present invention provides a carboxamide derivative of the formula (I'), its tautomeric or stereoisomeric form, or a salt thereof:



- 10 wherein m and n independently represent an integer from 0 to 2;

-X- represents -CH₂-CH₂-, -CH=CH-, or -C≡C-;

-R¹ represents -OR¹¹, -SR¹¹, -SOR¹¹, -SO₂R¹¹, -NR¹²R¹³, or -CHR¹⁴R¹⁵,

- 15 wherein R¹¹ represents (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl,

- 20 R¹² and R¹³ independently represent hydrogen, (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl, or R¹² and R¹³ together with the nitrogen atom to which they are attached, form a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

- 25 R¹⁴ and R¹⁵ independently represent hydrogen, (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkoxy optionally substituted by aryl or heteroaryl, or

R¹⁴ and R¹⁵ together with the CH to which they are attached, form a (C₃₋₈)cycloalkyl optionally interrupted by NH, or O, or R¹⁴ and R¹⁵ together with the CH to which they are attached, form a phenyl optionally substituted by hydroxy, halogen or (C₁₋₆) alkyl;

R² represents hydrogen, cyano, (C₁₋₆) alkoxy, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, (C₃₋₇)cycloalkyl, or (C₁₋₆) alkyl optionally substituted by amino, (C₁₋₆)alkylamino, or phenyl.

Yet another embodiment of the compounds of formula (I) or (I') are those wherein:

-R¹ represents -OR¹¹, -SR¹¹, -SOR¹¹, -SO₂R¹¹, -NR¹²R¹³, or -CHR¹⁴R¹⁵,

wherein R¹¹ represents (C₂₋₆)alkenyl substituted by aryl or heteroaryl, (C₂₋₆)alkynyl substituted by aryl or heteroaryl, or (C₁₋₆) alkyl substituted by aryl or heteroaryl;

R¹² and R¹³ independently represent (C₂₋₆)alkenyl substituted by aryl or heteroaryl, (C₂₋₆)alkynyl substituted by aryl or heteroaryl, or (C₁₋₆) alkyl substituted by aryl or heteroaryl;

R¹⁴ and R¹⁵ independently represent (C₂₋₆)alkenyl substituted by aryl or heteroaryl, (C₂₋₆)alkynyl substituted by aryl or heteroaryl, (C₁₋₆) alkyl substituted by aryl or heteroaryl, or (C₁₋₆) alkoxy substituted by aryl or heteroaryl.

Another embodiment of the compounds of formula (I) or (I') are those wherein: R¹ is phenoxy(C₁₋₆)alkyl, phenoxy(C₁₋₆)alkenyl, phenoxy(C₁₋₆)alkynyl, or phenyl(C₁₋₆)alkoxy.

Further embodiment of the compounds of formula (I) or (I') are those wherein R² is phenyl (C₁₋₆)alkyl.

Yet further embodiment of the compounds of formula (I) or (I') are those wherein R² is benzyl.

Further, the present invention provides a medicament which includes one of the compounds described above and optionally pharmaceutically acceptable excipient.

The Alkyl per se and "alk" and "alkyl" in alkoxy, alkanoyl, alkylamino, alkylaminocarbonyl, alkylaminosulphonyl, alkylsulphonylamino, alkoxycarbonyl, alkoxy-carbonylamino and alkanoylamino represent a linear or branched alkyl radical having generally 1 to 6, preferably 1 to 4 and particularly preferably 1 to 3 carbon atoms, representing illustratively and preferably methyl, ethyl, n-propyl, isopropyl, tert-butyl, n-pentyl and n-hexyl.

Alkoxy illustratively and preferably represents methoxy, ethoxy, n-propoxy, isopropoxy, tert-butoxy, n-pentoxy and n-hexoxy.

Alkylamino represents an alkylamino radical having one or two (independently selected) alkyl substituents, illustratively and preferably representing methylamino, ethylamino, n-propylamino, isopropylamino, tert-butylamino, n-pentylamino, n-hexyl-amino, N,N-dimethylamino, N,N-diethylamino, N-ethyl-N-methylamino, N-methyl-N-n-propylamino, N-isopropyl-N-n-propylamino, N-t-butyl-N-methylamino, N-ethyl-N-n-pentylamino and N-n-hexyl-N-methylamino.

Aryl per se represents a mono- to tricyclic aromatic carbocyclic radical having generally 6 to 14 carbon atoms, illustratively and preferably representing phenyl, naphthyl and phenanthrenyl.

Heteroaryl per se represents an aromatic mono- or bicyclic radical having generally 5 to 10 and preferably 5 or 6 ring atoms and up to 5 and preferably up to 4 hetero atoms selected from the group consisting of S, O and N, illustratively and preferably representing thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, pyridyl, pyrimidyl, pyridazinyl, indolyl, indazolyl, benzofuranyl, benzothiophenyl, quinolinyl, isoquinolinyl.

Hetero ring (heterocyclyl) per se represents a mono- or polycyclic, preferably mono- or bicyclic, nonaromatic heterocyclic radical having generally 4 to 10 and preferably 5 to 8 ring atoms and up to 3 and preferably up to 2 hetero atoms and/or hetero groups selected from the group consisting of N, O, S, SO and SO₂. The heterocyclyl radicals can be saturated or partially unsaturated. Preference is given to 5- to 8-membered monocyclic saturated heterocyclyl radicals having up to two hetero atoms selected from the group consisting of O, N and S, such as illustratively and preferably tetrahydrofuran-2-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrrolinyl, piperidinyl, morpholinyl, perhydroazepinyl.

EMBODIMENT OF THE INVENTION

The compound of the formula (I) of the present invention can be, but not limited to be, prepared by combining various known methods. In some embodiments, one or more of the substituents, such as amino group, carboxyl group, and hydroxyl group of the compounds used as starting materials or intermediates are advantageously protected by a protecting group known to those skilled in the art. Examples of the protecting groups are described in "Protective Groups in Organic Synthesis (3rd Edition)" by Greene and Wuts, John Wiley and Sons, New York 1999.

The compound of the formula (I) of the present invention can be, but not limited to be, prepared by the methods [A] below.

Method[A]




The compound of the formula (I) (wherein R^1 , R^2 , X, m, and n are the same as defined above, and Z represents C_{1-6} alkyl) or a salt thereof can be obtained by the hydrolysis of the starting material of formula (II).

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); alcohols such as methanol, ethanol, 1-propanol, isopropanol and tert-butanol, water, and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 100°C . The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.

The reaction can be advantageously carried out in the presence of a base including, for instance, an alkali metal alkoxide such as sodium methoxide, sodium ethoxide and potassium tert-butoxide; alkali metal hydroxide such as sodium hydroxide, lithium hydroxide and potassium hydroxide; and others.



The compound of formula (II) (wherein R^1 , R^2 , X, Z, m and n are the same as defined above) can be prepared by the reaction of compound (III) with amine (IV).

5

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

10

15


The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 0°C to 100°C . The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.

20

The reaction may be carried out using coupling agent including, for instance, carbodiimides such as N, N-dicyclohexylcarbodiimide and 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide, and others.

25

The compound (III) and amine (IV) can be commercially available or can be prepared by the use of known techniques.



Alternatively, the compound of formula (IIa) or (IIa)' (Wherein R², R¹¹, R¹², R¹³, X, Z, m and n are the same as defined above and A represents O or S) can be prepared by the reaction of compound (V) or (V)' with reagents (VI) or (VI)', respectively (wherein Y represents a leaving group, e.g., halogen and alkylsulfonic acid), such as benzyl chloride or benzyl bromide.

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); ketones such as acetone; alcohols such as methanol, ethanol, 1-propanol, isopropanol and tert-butanol, and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 0°C to 100°C. The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.

The reaction can be advantageously carried out in the presence of a base including, for instance, an alkali metal hydride such as sodium hydride or potassium hydride; alkali metal alkoxide such as sodium methoxide, sodium ethoxide and potassium tert-butoxide; alkali metal hydroxide such as sodium hydroxide and potassium hydroxide; alkali metal carbonates such as sodium carbonate and potassium carbonate; alkali metal hydrogen carbonates such as sodium hydrogen carbonate and potassium hydrogen carbonate; alkaline earth metal alkoxides such as magnesium ethoxide; organic amines such as pyridine, triethylamine and N,N-diisopropylethylamine, dimethylaniline, diethylaniline and others.

The compound (V) or (V)' can be commercially available or can be prepared by either the use of the similar procedure for the preparation of the compound of formula (II) or known techniques. The compound (VI) or (VI)' can be commercially available or can be prepared by the use of known techniques.

When the compound shown by the formula (I) or a salt thereof has an asymmetric carbon in the structure, their optically active compounds and racemic mixtures are also included in the scope of the present invention.

Typical salts of the compound shown by the formula (I) include salts prepared by reaction of the compounds of the present invention with a mineral or organic acid, or an organic or inorganic base. Such salts are known as acid addition and base addition salts, successively.

Acids to form salts include inorganic acids such as, without limitation, sulfuric acid, phosphoric acid, hydrochloric acid, hydrobromic acid, hydriodic acid and the like, and organic acids, such as, without limitation, p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromophenylsulfonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like.

Base addition salts include those derived from inorganic bases, such as, without limitation, ammonium hydroxide, alkaline metal hydroxide, alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, and organic bases, such as, without limitation, ethanolamine, triethylamine, tris(hydroxymethyl)aminomethane, and the like. Examples of inorganic bases include, sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like.

The compound of the present invention or a salts thereof, depending on its substituents, may be modified to form lower alkylesters or known other esters; and/or hydrates or other solvates. Those esters, hydrates, and solvates are included in the scope of the present invention.

The compound of the present invention may be administered in oral forms, such as, without limitation normal and enteric coated tablets, capsules, pills, powders, granules, elixirs, tinctures, solution, suspensions, syrups, solid and liquid aerosols and emulsions. They may also be administered in parenteral forms, such as, without limitation, intravenous, intraperitoneal, subcutaneous, intramuscular, and the like forms, well-known to those of ordinary skill in the pharmaceutical arts. The compounds of the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal delivery systems well-known to those of ordinary skilled in the art.

The dosage regimen with the use of the compounds of the present invention is selected by one of ordinary skill in the arts, in view of a variety of factors, including, without limitation, age, weight, sex, and medical condition of the recipient, the severity of the condition to be treated, the route of administration, the level of metabolic and excretory function of the recipient, the dosage form employed, the particular compound and salt thereof employed.

The compounds of the present invention are preferably formulated prior to administration together with one or more pharmaceutically-acceptable excipients. Excipients are inert substances such as, without limitation carriers, diluents, flavoring agents, sweeteners, lubricants, solubilizers, suspending agents, binders, tablet
5 disintegrating agents and encapsulating material.

Yet another embodiment of the present invention is pharmaceutical formulation comprising a compound of the invention and one or more pharmaceutically-acceptable excipients that are compatible with the other ingredients of the
10 formulation and not deleterious to the recipient thereof. Pharmaceutical formulations of the invention are prepared by combining a therapeutically effective amount of the compounds of the invention together with one or more pharmaceutically-acceptable excipients. In making the compositions of the present invention, the active
15 ingredient may be mixed with a diluent, or enclosed within a carrier, which may be in the form of a capsule, sachet, paper, or other container. The carrier may serve as a diluent, which may be solid, semi-solid, or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions, solutions, syrups, aerosols, ointments, containing, for example, up to 10%
20 by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

For oral administration, the active ingredient may be combined with an oral, and non-toxic, pharmaceutically-acceptable carrier, such as, without limitation, lactose, starch, sucrose, glucose, sodium carbonate, mannitol, sorbitol, calcium carbonate,
25 calcium phosphate, calcium sulfate, methyl cellulose, and the like; together with, optionally, disintegrating agents, such as, without limitation, maize, starch, methyl cellulose, agar bentonite, xanthan gum, alginic acid, and the like; and optionally, binding agents, for example, without limitation, gelatin, natural sugars, beta-lactose, corn sweeteners, natural and synthetic gums, acacia, tragacanth, sodium alginate,
30 carboxymethylcellulose, polyethylene glycol, waxes, and the like; and, optionally, lubricating agents. For example, without limitation, magnesium stearate, sodium

stearate, stearic acid, sodium oleate, sodium benzoate, sodium acetate, sodium chloride, talc, and the like.

5 In powder forms, the carrier may be a finely divided solid which is in admixture with the finely divided active ingredient. The active ingredient may be mixed with a carrier having binding properties in suitable proportions and compacted in the shape and size desired to produce tablets. The powders and tablets preferably contain from about 1 to about 99 weight percent of the active ingredient which is the novel composition of the present invention. Suitable solid carriers are magnesium
10 carboxymethyl cellulose, low melting waxes, and cocoa butter.

Sterile liquid formulations include suspensions, emulsions, syrups and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent, or a mixture of both sterile water
15 and sterile organic solvent.

The active ingredient can also be dissolved in a suitable organic solvent, for example, aqueous propylene glycol. Other compositions can be made by dispersing the finely divided active ingredient in aqueous starch or sodium carboxymethyl cellulose
20 solution or in a suitable oil.

The formulation may be in unit dosage form, which is a physically discrete unit containing a unit dose, suitable for administration in human or other mammals. A unit dosage form can be a capsule or tablets, or a number of capsules or tablets. A
25 "unit dose" is a predetermined quantity of the active compound of the present invention, calculated to produce the desired therapeutic effect, in association with one or more excipients. The quantity of active ingredient in a unit dose may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved.

Typical oral dosages of the present invention, when used for the indicated effects, will range from about 0.01mg /kg/day to about 100 mg/kg/day, preferably from 0.1 mg/kg/day to 30 mg/kg/day, and most preferably from about 0.5 mg/kg/day to about 10 mg/kg/day. In the case of parenteral administration, it has generally proven advantageous to administer quantities of about 0.001 to 100mg /kg/day, preferably from 0.01 mg/kg/day to 1 mg/kg/day. The compounds of the present invention may be administered in a single daily dose, or the total daily dose may be administered in divided doses, two, three, or more times per day. Where delivery is via transdermal forms, of course, administration is continuous:

Examples

The present invention will be described in detail below in the form of examples, but they should by no means be construed as defining the metes and bounds of the present invention.

In the examples below, all quantitative data, if not stated otherwise, relate to percentages by weight.

Melting points are uncorrected. Liquid Chromatography - Mass spectroscopy (LC-MS) data were recorded on a Micromass Platform LC with Shimadzu Phenomenex ODS column(4.6 mm X 30 mm) flushing a mixture of acetonitrile-water (9:1 to 1:9) at 1 ml/min of the flow rate. Mass spectra were obtained using electrospray (ES) ionization techniques (micromass Platform LC). TLC was performed on a precoated silica gel plate (Merck silica gel 60 F-254). Silica gel (WAKO-gel C-200 (75-150 μ m)) was used for all column chromatography separations. All chemicals were reagent grade and were purchased from Sigma-Aldrich, Wako pure chemical industries, Ltd., Tokyo kasei kogyo Co., Ltd., Nacalai tesque, Inc., Watanabe Chemical Ind. Ltd., Maybridge plc, Lancaster Synthesis Ltd., Merck KGaA, Kanto Chemical Co., Ltd.

The effect of the present compounds were examined by the following assays and pharmacological tests.

[Measurement of the [3 H]-iloprost binding to HEL cells] (Assay 1)

A human erythroleukemia cell line, HEL 92.1.7, was purchased from American Type Culture Collection and maintained in RPMI-1640 medium (Gibco BRL) supplemented with 10% fetal calf serum (FCS), 2 mM glutamine, 4.5 g/L glucose, 10 mM Hepes, 1 mM sodium pyruvate, 100 U/ml penicillin, and 100 μ g/ml streptomycin in a humidified 5% CO₂ atmosphere at 37°C. Cells were collected with

centrifugation and washed with binding assay buffer (BAB: 50 mM Tris-HCl, 5 mM MgCl₂ (pH 7.5)). Cells were suspended at the density of 6.25×10^6 cells/ml in BAB, and one million cells in 160 μ l aliquot of cell suspension were put in a well of 96 well plate (Falcon). Then, 20 μ l of compound solution, 100 μ M of iloprost (for non-specific binding), or buffer alone (total binding), diluted with 1% DMSO in BAB was added. Finally, another 20 μ l containing [³H]-iloprost (0.02 μ Ci, 0.5-1 pmol) in BAB was added and incubated at room temperature for 30 min with a gentle shaking. Cell suspension was then transferred to a well of MultiScreen plate with GF/C glass filters (Millipore) to harvest cells. Cells were washed twice with 200 μ l of ice-cold BAB and the plate was kept at 55°C for 30 min to dry filters. The filter in the well was punched out to a counting tube and 2 ml of Ultima Gold XR (Packard) was added. [³H]-radio activity in the filter was measured by a liquid scintillation counter (Beckman).

[Iloprost-induced cAMP production assay in HEL cells] (Assay 2)

HEL cells were collected with centrifugation and washed with cAMP assay buffer (CAB: Hank's balanced salt solution, 17 mM Hepes, 0.1% bovine serum albumin, 1 mM IBMX, 0.4% DMSO, and 1 mM L-ascorbic acid sodium salt (pH 7.4)). Cells were suspended at the density of 2.5×10^5 cells/ml in CAB, and twenty thousand cells in 80 μ l aliquot of cell suspension were put in a well of 96 well plate (Falcon). Then, 10 μ l of compound solution diluted with 1% DMSO in CAB or buffer alone was added. The plate was incubated at 37°C for 30 min. Then, another 10 μ l containing 100 nM iloprost in CAB or buffer alone was added and further incubated at 37°C for 30 min. cAMP content in the well was measured by a cAMP ELISA kit (Applied Biosystems).

[Measurement of rhythmic bladder contraction in anesthetized rats]**(1) Animals**

Female Sprague-Dawley rats (200~250 g / Charles River Japan) were used.

(2) Rhythmic bladder contraction in anesthetized rats

Rats were anesthetized by intraperitoneal administration of urethane (Sigma) at 1.25 g/kg. The trachea was cannulated with a polyethylene tube (HIBIKI, No.8) to facilitate respiration; and a cannula (BECTON DICKINSON, PE-50) was placed in the left femoral vein for intravenous administration of testing compounds. The abdomen was opened through a midline incision, and after both ureters were cut, a water-filled balloon (about 1 ml capacity) was inserted through the apex of the bladder dome. The balloon was connected to a pressure transducer onto a polygraph. Rhythmic bladder contraction was elicited by raising up intravesical pressure to approximately 15 cm H₂O. After the rhythmic bladder contraction was stable, a testing compound was administered intravenously. Activity was estimated by measuring disappearance time and amplitude of the rhythmic bladder contraction. The effect on amplitude of bladder contractions was expressed as a percent suppression of the amplitude of those after the disappearance was recovered. Experimental values were expressed as the mean±S.E.M. The testing compounds-mediated inhibition of the rhythmic bladder contraction was evaluated using Student's t-test. A probability level less than 5% was accepted as significant difference.


Results of PGI₂ kinase assay is shown in Examples and tables of the Examples below. The data corresponds to the compounds as yielded by solid phase synthesis and thus to levels of purity of about 40 to 90%. For practical reasons, the compounds are grouped in three classes of activity as follows:

IC₅₀ = A 1 μM < B 10 μM < C

The compounds of the present invention also show excellent selectivity, and strong activity in in vivo assays.


Example 1:

(1) 4-Chloromethylbenzyl Alcohol



To a solution of 4-chloro-4-toluic acid in tetrahydrofuran (THF, 60 ml) was added 1 M borane THF solution (90 ml). The mixture was stirred at room temperature overnight and quenched by addition of methanol (50 ml). The solvent was evaporated off and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1 to 3/1) to obtain 4-chloromethylbenzyl Alcohol (8.84 g, 96%) as a colorless solid.

(2) 4-Phenoxymethylbenzyl Alcohol




A mixture of 4-chloromethylbenzyl alcohol (0.80 g), phenol (0.48 g), 85% potassium hydroxide (0.76 g) and dimethylsulfoxide (DMSO, 15 ml) was stirred at room temperature overnight and poured into a mixture of water (50 ml) and ethyl acetate (50 ml). The organic layer was washed with brine and dried over sodium sulfate. The solvent was removed off and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 3/1) to obtain 4-phenoxymethylbenzyl alcohol (0.83 g, 76%) as colorless granules.

(3) 4-Phenoxymethylbenzaldehyde




To a solution of Dess-Martin reagent (1.79 g) in dichloromethane (10 ml) was dropwise added a solution of 4-phenoxymethylbenzyl alcohol (0.82 g) at room temperature. The mixture was stirred at room temperature for 30 min and poured into 1N NaOH water solution (30 ml). The organic layer was washed with water and dried over sodium sulfate. The solvent was removed off and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to obtain 4-phenoxymethylbenzaldehyde (0.67 g, 81%) as a colorless solid.

(4) tert-Butyl 4-Phenoxymethylcinnamate




To a solution of tert-butyl diethoxyphosphorylacetate (0.66 g) in THF (10 ml) was added 60% sodium hydride (0.10 g) at 0°C. The mixture was stirred for 1 hr on an ice-water bath and a solution of 4-phenoxymethylbenzaldehyde in THF (1 ml) was added dropwise. The reaction mixture was stirred at room temperature overnight and poured into saturated ammonium chloride water solution (50 ml). The resulting suspension was extracted with ethyl acetate and the organic layer was washed dried over sodium sulfate. The solvent was removed and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to obtain tert-butyl 4-phenoxymethylcinnamate (0.73 g, 100%) as a colorless solid.

(5) N-(4-Phenoxymethylcinnamoyl)phenylalanine Methyl Ester



5 A mixture of tert-butyl 4-phenoxymethylcinnamate (0.20 g), trifluoroacetic acid (TFA, 1 ml) and dichloromethane (1 ml) was allowed to stand for 2.5 hr at room temperature. The solvent was removed in vacuo and the residue was dissolved in N,N-dimethylformamide (DMF, 5 ml). To the solution were added phenylalanine methyl ester (0.15 g), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI, 0.18 g), 1-hydroxybenzotriazole (HOBt, 0.12 g) and triethylamine (0.12 ml). The mixture was stirred at room temperature overnight and poured into a mixture of water (30 ml) and ethyl acetate (20 ml). The organic layer was washed with water and dried over sodium sulfate. The solvent was removed off and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to obtain N-(4-phenoxymethylcinnamoyl)phenylalanine methyl ester (0.22 g, 88%) as a colorless solid.

(6) N-(4-phenoxymethylcinnamoyl)phenylalanine



20 To a solution of N-(4-phenoxymethylcinnamoyl)phenylalanine methyl ester (70 mg) in methanol (2 ml) was added 1N lithium hydroxide water solution (0.2 ml). The mixture was stirred at 50°C overnight and concentrated in vacuo. The residue was

dissolved in water and acidified with 1N hydrochloric acid. The resulting suspension was extracted with ethyl acetate. The organic layer was washed with brine and dried over sodium sulfate. The solvent was removed off and the residue was triturated with diisopropyl ether to obtain N-(4-phenoxyethylcinnamoyl)phenylalanine (59 mg, 86%) as a colorless solid.


mp 205 °C; Calcd [M+1]: 402, Found: m/z 402.

Molecular weight: 401.47

Activity grade assay 1: A


Example 2:

(1) tert-Butyl 3-(4-Phenoxyethylphenyl)propionate



To a mixture of tert-butyl 4-phenoxyethylcinnamate (see: example 1-(4), 0.20 g) and nickel chloride hexahydrate (0.02 g) in methanol (4 ml) was added sodium borontetrahydride (0.05 g) on an ice-water bath. The mixture was stirred at room temperature for 1 hr and quenched with saturated ammonium chloride water solution. The resulting suspension was extracted with ethyl acetate and the organic layer was washed with water and dried over sodium sulfate. The solvent was removed off and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 4/1) to obtain tert-butyl 3-(4-phenoxyethylphenyl)propionate (0.168 g, 84%) as a colorless solid.

(2) N-[3-(4-Phenoxymethylphenyl)propionyl]phenylalanine



5

To a solution of tert-butyl 3-(4-phenoxymethylphenyl)propionate (0.10 g) in ethanol (2 ml) was added 1N lithium hydroxide water solution (0.7 ml). The mixture was stirred at 60°C for 3 hr and concentrated in vacuo. The residue was suspended in a mixture of 1N hydrochloric acid (0.7 ml), water (5 ml) and ethyl acetate (10 ml) and the organic layer was washed with brine and dried over sodium sulfate. The solvent was removed off and the residue was used for the following steps towards N-[3-(4-Phenoxymethylphenyl)propionyl]phenylalanine, according to the procedures for the synthesis of N-(4-phenoxymethylcinnamoyl)phenylalanine (See: example 1-(5) and (6)).

15

mp 173-174 °C; Calcd [M+1]: 404, Found: m/z 404.


Molecular weight: 403.48

Activity grade assay 1: A

20

Example 3:


(1) 1-Iodo-4-(phenoxymethyl)benzene



25


A mixture of 4-iodobenzyl bromide (1 g), phenol (0.286 g), potassium carbonate (0.530 g) and DMF (20 ml) was stirred at room temperature overnight. The volatiles were removed off in vacuo and the residue was suspended in a mixture of ethyl acetate and water. The organic layer was separated to be washed with brine and dried over sodium sulfate. The solvent was removed and the residue was purified by silica gel column chromatography to obtain 1-iodo-4-(phenoxy)methylbenzene (0.918 g, 93%) as pale yellow flakes.

(2) Methyl 4-Phenoxyethylphenylpropiolate



To a solution of 1-iodo-4-(phenoxy)methylbenzene (0.40 g) and methyl propiolate (0.43 g) in THF (8 ml) were added Biskis(triphenylphosphine)palladium dichloride (18 mg), cuprous iodide (10 mg) and potassium carbonate (0.36 g). The mixture was stirred at 80°C and concentrated in vacuo. The residue was purified by silica gel column chromatography (hexane/ethyl acetate = 10/1) to obtain methyl 4-phenoxyethylphenylpropiolate (0.155 g, 45%) as colorless flakes.

(3) N-(4-Phenoxyethylphenylpropioloyl)phenylalanine



According to the procedure for the synthesis of N-(4-phenoxy-methyl-cinnamoyl)phenylalanine (See: example 1-(5) and (6)) from tert-butyl 4-phenoxy-methylcinnamate, N-(4-phenoxy-methylphenylpropioloyl)phenylalanine was prepared from 4-phenoxy-methylphenylpropionic acid, which was obtained from the
5 corresponding methyl ester by the hydrolysis with 1N lithium hydroxide in ethanol:

mp 146 °C; Calcd [M+1]: 400, Found: m/z 400.


Molecular weight: 399.44

Activity grade assay 1: A

10

Example 4:

(1) 4-Benzyloxybenzaldehyde




15

20

To a solution of 4-hydroxybenzaldehyde (1.00 g) in DMF (30 ml) were added benzyl chloride (1.24 g) and potassium carbonate (1.36 g). The mixture was stirred at room temperature overnight and at 60°C for 2 hr. The reaction mixture was concentrated in vacuo and the residue was suspended in a mixture of ethyl acetate and water. The organic layer was separated to be washed with brine and dried over magnesium sulfate. The solvent was removed in vacuo and the residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to obtain 4-benzyloxybenzaldehyde (1.82 g, 100%) as a colorless solid.

25

(2) N-(4-Benzyloxycinnamoyl)phenylalanine



5

According to the procedures for the synthesis of tert-butyl 4-phenoxy-methylcinnamate (See: example 1-(4)), 4-benzyloxybenzaldehyde was subjected to the described Horner-Emmons reaction to obtain tert-butyl 4-benzyloxycinnamate, followed by hydrolysis with lithium hydroxide in ethanol using the procedure described in example 2-(2). Resulting 4-benzyloxycinnamic acid was coupled with phenylalanine methyl ester and hydrolyzed with lithium hydroxide in ethanol to obtain N-(4-benzyloxycinnamoyl)phenylalanine, according to the procedure for the synthesis of N-(4-phenoxy-methylcinnamoyl)phenylalanine (See: example 1-(6)).

10

15


mp 220 °C; Calcd [M+1]: 402, Found: m/z 402.

Molecular weight: 401.46

Activity grade assay 1: A

Claims

1. A carboxamide derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof:



wherein m and n independently represent an integer from 0 to 2;

-X- represents $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$, or $-\text{C}\equiv\text{C}-$;

-R¹ represents $-\text{OR}^{11}$, $-\text{SR}^{11}$, $-\text{SOR}^{11}$, $-\text{SO}_2\text{R}^{11}$, $-\text{NR}^{12}\text{R}^{13}$, or $-\text{CHR}^{14}\text{R}^{15}$,

wherein R^{11} represents (C_{2-6}) alkenyl optionally substituted by aryl or heteroaryl, (C_{2-6}) alkynyl optionally substituted by aryl or heteroaryl, (C_{1-6}) alkyl optionally substituted by aryl or heteroaryl,

R^{12} and R^{13} independently represent hydrogen, (C_{2-6}) alkenyl optionally substituted by aryl or heteroaryl, (C_{2-6}) alkynyl optionally substituted by aryl or heteroaryl, (C_{1-6}) alkyl optionally substituted by aryl or heteroaryl, or R^{12} and R^{13} together with the nitrogen atom to which they are attached, form a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

R^{14} and R^{15} independently represent hydrogen, (C_{2-6}) alkenyl optionally substituted by aryl or heteroaryl, (C_{2-6}) alkynyl optionally substituted by aryl or heteroaryl, (C_{1-6}) alkyl optionally substituted by aryl or heteroaryl, (C_{1-6}) alkoxy optionally substituted by aryl or heteroaryl, or


R^{14} and R^{15} together with the CH to which they are attached, form a (C₃₋₈)cycloalkyl optionally interrupted by NH, or O, or R^{14} and R^{15} together with the CH to which they are attached, form a phenyl optionally substituted by hydroxy, halogen or (C₁₋₆) alkyl;

5

R^2 represents hydrogen, cyano, (C₁₋₆) alkoxy, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, (C₃₋₇)cycloalkyl, or (C₁₋₆) alkyl optionally substituted by amino, (C₁₋₆)alkylamino, or phenyl.

10

2. A carboxamide derivative of the formula (I'), its tautomeric or stereoisomeric form, or a salt thereof:



15

wherein m and n independently represent an integer from 0 to 2;

-X- represents -CH₂-CH₂-, -CH=CH-, or -C≡C-;

-R¹ represents -OR¹¹, -SR¹¹, -SOR¹¹, -SO₂R¹¹, -NR¹²R¹³, or -CHR¹⁴R¹⁵,

20

wherein R¹¹ represents (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl,

25

R^{12} and R^{13} independently represent hydrogen, (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl, or R^{12} and R^{13} together with the nitrogen atom to which

they are attached, form a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

5 R^{14} and R^{15} independently represent hydrogen, (C₂₋₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂₋₆)alkynyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkyl optionally substituted by aryl or heteroaryl, (C₁₋₆) alkoxy optionally substituted by aryl or heteroaryl, or

10 R^{14} and R^{15} together with the CH to which they are attached, form a (C₃₋₈)cycloalkyl optionally interrupted by NH, or O, or R^{14} and R^{15} together with the CH to which they are attached, form a phenyl optionally substituted by hydroxy, halogen or (C₁₋₆) alkyl;

15 R^2 represents hydrogen, cyano, (C₁₋₆) alkoxy, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, (C₃₋₇)cycloalkyl, or (C₁₋₆) alkyl optionally substituted by amino, (C₁₋₆)alkylamino, or phenyl.

20 3. The carboxamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1 or 2,

 wherein $-R^1$ represents $-OR^{11}$, $-SR^{11}$, $-SOR^{11}$, $-SO_2R^{11}$, $-NR^{12}R^{13}$, or $-CHR^{14}R^{15}$,

25 wherein R^{11} represents (C₂₋₆)alkenyl substituted by aryl or heteroaryl, (C₂₋₆)alkynyl substituted by aryl or heteroaryl, or (C₁₋₆) alkyl substituted by aryl or heteroaryl;

30 R^{12} and R^{13} independently represent (C₂₋₆)alkenyl substituted by aryl or heteroaryl, (C₂₋₆)alkynyl substituted by aryl or heteroaryl, or (C₁₋₆) alkyl substituted by aryl or heteroaryl;

R¹⁴ and R¹⁵ independently represent (C₂₋₆)alkenyl substituted by aryl or heteroaryl, (C₂₋₆)alkynyl substituted by aryl or heteroaryl, (C₁₋₆) alkyl substituted by aryl or heteroaryl, or (C₁₋₆) alkoxy substituted by aryl or heteroaryl.

- 5
4. The carboxamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1 or 2,

10 wherein R¹ is phenoxy(C₁₋₆)alkyl, phenoxy(C₁₋₆)alkenyl, phenoxy(C₁₋₆)alkynyl, or phenyl(C₁₋₆)alkoxy.

- 15
5. The carboxamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1 or 2,

wherein R² is phenyl (C₁₋₆)alkyl.

- 20
6. The carboxamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1 or 2,

wherein R² is benzyl.

- 25
7. A medicament comprising the carboxamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 or 2 as an active ingredient.

8. The medicament as claimed in claim 7, further comprising one or more pharmaceutically acceptable excipients.

9. The medicament as claimed in claim 7, wherein the carboxamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof is a PGI2 antagonist.
- 5 10. An agent to treat or prevent a urological disorder or disease; comprising the carboxamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 as an active ingredient.
- 10 11. An agent to treat pain; comprising the carboxamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 as an active ingredient.
- 15 12. An agent to treat hypotension; comprising the carboxamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 as an active ingredient.
- 20 13. An agent to treat hemophilia and hemorrhage; comprising the carboxamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 as an active ingredient.
- 25 14. An agent to treat inflammation; comprising the carboxamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 as an active ingredient.
- 30 15. Use of compounds according to any of Claims 1 to 3 for manufacturing a medicament for the treatment and/or prophylaxis of urological disorders.
16. Process for controlling urological disorders in humans and animals by administration of a PGI2-antagonistically effective amount of at least one compound according to any of Claims 1 to 3.

Carboxamides derivatives

ABSTRACT

The present invention relates to carboxamides which are useful as an active ingredient of pharmaceutical preparations. The carboxamides of the present invention have PGI₂ antagonistic activity, and can be used for the prophylaxis and treatment of diseases associated with PGI₂ activity.

Such diseases include urological diseases or disorder as follows: bladder outlet obstruction, overactive bladder, urinary incontinence, detrusor hyper-reflexia, detrusor instability, reduced bladder capacity, frequency of micturition, urge incontinence, stress incontinence, bladder hyperreactivity, benign prostatic hypertrophy (BPH), prostatitis, urinary frequency, nocturia, urinary urgency, pelvic hypersensitivity, urethritis, pelvic pain syndrome, prostatodynia, cystitis, or idiopathic bladder hypersensitivity.

The compounds of the present invention are also useful for treatment of pain including, but not limited to inflammatory pain, neuropathic pain, acute pain, chronic pain, dental pain, premenstrual pain, visceral pain, headaches, and the like; hypotension; hemophilia and hemorrhage; and inflammation, since the diseases also relate to PGI₂.